

The Henry Ford Study

A 5 years hidden study of vaccinated versus unvaccinated children comes out with shocking results

A few weeks ago, a new study of vaccinated versus unvaccinated children came out. By its sheer size, this study cripples all previous vaxxed/un-vaxxed studies, and it clearly deserves all attention.

By **Ivo Zvardon**

October 7, 2025



It all started in 2020, when the Institute of Medicine (IOM) ordered the Division of Infectious Diseases of the Henry Ford hospitals a study of vaccinated versus fully unvaccinated children. The goal was to finally conduct an extensive study – since a fully comprehensive, data-based study of this magnitude has never been performed.

The purpose was to come to results positive for the pharmaceutical industry – that vaccines do not cause health conditions in vaccinated kids. As the authors of the paper mention (Study, p.3):

Addressing this significant data gap could allay parental concerns and bolster vaccine confidence.

Unfortunately for the pharma companies, the results that came out of this study are the exact opposite – they demonstrate that unvaccinated children are many times healthier than those who got one or more jabs. In view of this conclusion, the study was swept under the rug and kept secret for years.

Aaron Siri, an attorney who specializes in the field of civil litigation, came up with the study and, on September 9, testified in a hearing at the Senate.

The study never underwent peer-review, since the authors – lead author Marcus Zervos, MD, Amy Tang PhD, Agigail Chatfield, MS, Lois Lamerato, PhD – said they would have lost their jobs had they decided to make it public. So, hidden it stayed until last month.

What makes the Henry Ford study so important?

The large number of participants. 18 468 children took part in the study.

The study's scientific robustness. Each participant was enrolled in the Henry Ford health insurance plan, so the data of all the participants – registries from the Henry Ford Health System (HFHS), the Health Alliance Plan (HAP) and State of Michigan immunization registry - are trackable.

Its scale. It is the first study of this size, conducted by health authorities, that compares fully unvaccinated with vaccinated children (fully or partially). Most previous studies conducted by health agencies were comparing, for example, unvaccinated children with children that got a single specific immunization, or children that underwent one or more specific vaccines with children fully vaccinated.

The study's credibility – performed by one of the largest health institutions in the U.S. As the study authors mention (Study, p.4):

The study was reviewed and approved by HFHS's Institutional Review Board and conducted in accordance with the International Society for Pharmacoepidemiology's Guidelines for Good Pharmacoepidemiology Practices
(https://www.pharmacoepi.org/resources/guidelines_08027.cfm)

Study design

The study included children born between 2000 and 2016. Subjects were observed from birth until December 2017.

The authors divided the participants into two groups:

- the fully unvaccinated (1 957 children)
- those, who received at least one vaccine during their enrollment in the plan (16 511 children).

Results of the study

Vaccinated children were:

- **6.16 times more likely to develop an autoimmune disease**
- **4.09 times more likely to develop asthma**
- **6.15 times more likely to develop a neurodevelopmental disorder (ADHD, Behavioral disability, Developmental delay, Learning disability, Intellectual disability, Speech disorder, Motor disability, tics).**
- **2.64 times more likely to develop an atopic disease (a group of allergic conditions)**

No significant connection between vaccination and cancer or food allergies was found.

See chart below (study p.17) (The 3rd column shows the odds of developing a certain health condition, adjusted to the number of participants in each group):

Table 2. Incidence of Chronic Health Conditions Stratified by Vaccine Exposure Status*

Outcome	Any Vaccine Exposure	No Vaccine Exposure	IRR (95% CI)	P
	N (Incidence per 1,000,000 pt-yrs)	N (Incidence per 1,000,000 pt-yrs)		
Chronic Health Condition	4,732 (277.3)	160 (111.7)	2.48 (2.12-2.91)	<0.0001
Asthma	2,867 (145.6)	52 (35.6)	4.09 (3.11-5.38)	<0.0001
Atopic Disease	946 (41.2)	23 (15.6)	2.64 (1.74-3.99)	<0.0001
Autoimmune Disease	201 (8.4)	2 (1.4)	6.16 (1.53-24.79)	0.01
Brain Dysfunction	8 (0.3)	0 (0.0)	∞	
Cancer	169 (7.0)	13 (8.8)	0.79 (0.45-1.39)	0.42
Diabetes	42 (1.7)	0 (0.0)	∞	
Food Allergy	577 (24.3)	30 (20.5)	1.19 (0.82-1.71)	0.36
Mental Health Disorder	341 (15.9)	5 (4.5)	3.50 (1.45-8.46)	<0.01
Neurodevelopmental Disorder	1,029 (50.2)	9 (8.2)	6.15 (3.19-11.86)	<0.0001
ADHD	262 (12.1)	0 (0.0)	∞	
Autism	23 (1.1)	1 (0.9)	1.16 (0.16-8.62)	0.88
Behavioral Disability	165 (7.6)	0 (0.0)	∞	
Developmental Delay	219 (10.1)	3 (2.7)	3.74 (1.20-11.68)	0.02
Learning Disability	65 (3.0)	0 (0.0)	∞	
Intellectual Disability	5 (0.2)	0 (0.0)	∞	
Speech Disorder	463 (21.8)	6 (5.4)	4.02 (1.80-9.00)	<0.001
Motor Disability	150 (6.9)	2 (1.8)	3.83 (0.95-15.47)	0.06
Tics	46 (2.1)	0 (0.0)	∞	
Other Psychological Disability	9 (0.4)	0 (0.0)	∞	
Neurological Disorder	127 (5.2)	12 (8.1)	0.64 (0.35-1.16)	0.14
Seizure Disorder	319 (13.3)	12 (8.2)	1.63 (0.92-2.91)	0.09

* Incident rate ratios could not be calculated for brain dysfunction, diabetes, ADHD, tics, or behavioral, learning, intellectual, or other psychological disability since all cases occurred in the group exposed to vaccination and no cases occurred in the unexposed group.

Interpretation

It seems that, due to certain biologic mechanisms, certain individuals are at greater risk to develop a health condition after vaccination than others. It is believed that the antigens contained in vaccines trigger a specific immune response, with some recipients more susceptible to immunological effects. As the authors of the study mention (Study, p.11):

The results of this study, while preliminary, suggest that we currently underestimate the group susceptible to an adverse vaccine effect.


Final conclusions of the study:

In total, vaccinated children were 2.48 times more likely to develop a chronic health condition than unvaccinated children.

The overall probability of being free of a chronic health condition at 10 years of follow-up was 43 percent in the group exposed to vaccination, and 83 percent in the unexposed group.

In other words, on average 1 out of 2 vaccinated children will develop a health condition over a 10-year period, while among the unvaccinated ones, only 1 out of five will develop a health condition. (Chart p.19 of the study):

Figure 1. Kaplan Meier Curve: 10-year Chronic Disease-Free Survival by Vaccine Exposure



The authors conclude (study p.15):

Conclusion: *In this study, we found vaccine exposure in children was associated with an increased risk of developing a chronic health disorder. This association was primarily driven by increased risk for asthma, atopy, eczema, autoimmune disease and neurodevelopmental disorders. This suggests that in certain susceptible children, exposure to vaccination may increase the likelihood of developing a chronic health condition, particularly for one of these conditions. Our preliminary findings cannot prove causality and warrant further investigation.*

Sen. Ron Johnson (R-Wis.), subcommittee chair who organized the hearing at Congress – which was the 3rd hearing about vaccine safety this year - said he hoped the hearing would open people’s minds so that “more Americans have their eyes open to the reality and truth.”

I urge everybody to read this important study – it is about 20 pages and can be found here:
<https://www.documentcloud.org/documents/26089210-henry-ford-vaccinated-unvaccinated-study/>

The conclusions of this paper are in many ways similar to the results of other previous vaxxed/un-vaxxed studies (all of them were, however, smaller in scale):

- Hooker and Miller (Hooker and Miller, 2020) found a 4.49 higher risk of asthma and a 2.18 times higher risk of development delays in the vaccinated category, compared to the unvaccinated one. (1)
- Mawson (Mawson et al., 2017) found a 2.7 higher risk of developing a neurodevelopmental disorder for kids that had been vaccinated, compared to the unvaccinated. (2)

Final thoughts

The pharmaceutical companies, the medias on their payrolls and our health authorities have a long tradition of claiming that robust studies of vaccinated versus unvaccinated children do not exist – creating a loophole allowing them to carry on with the assertion that vaccines are safe.

Considering the scale, the scientific rigorousness and robustness of the Henry Ford study, it is obvious that the industry cannot claim any longer that children’s immunizations are safe. In view of the alarming outcomes of this study, it is clear as day that proper and honest safety testing of vaccines must finally be performed.

Until then, the vaccination of our children – in many countries’ compulsory under the vaccination schedule – needs to be suspended.

Ivo Zvardon

ivozvardon@gmail.com
@ivozvardon

Annex 1

Analysis of health outcomes in vaccinated and unvaccinated children: Developmental delays, asthma, ear infections and gastrointestinal disorders

<https://journals.sagepub.com/doi/full/10.1177/2050312120925344>

Annex 2

Preterm birth, vaccination and neurodevelopmental disorders: a cross-sectional study of 6 to 12-year-old vaccinated and unvaccinated children

<https://www.oatext.com/Preterm-birth-vaccination-and-neurodevelopmental-disorders-a-cross-sectional-study-of-6-to-12-year-old-vaccinated-and-unvaccinated-children.php>